

The University of Alabama
January 20, 2017

Stronger Rolle's Theorem for Complex Polynomials

Hristo Sendov

Department of Statistical and Actuarial Sciences
The University of Western Ontario

London, Ontario
Canada N6A 5B7
hssendov@stats.uwo.ca

Every Calculus student is familiar with the classical Rolle's theorem stating that if a real polynomial p satisfies $p(-1) = p(1)$, then it has a critical point in $(-1, 1)$. In 1934, L. Tschakaloff strengthened this result by finding a *minimal* interval, contained in $(-1, 1)$, that holds a critical point of every real polynomial with $p(-1) = p(1)$, up to a fixed degree. In 1936, he expressed a desire to find an analogue of his result for complex polynomials.

This talk will present the following Rolle's theorem for complex polynomials. If $p(z)$ is a complex polynomial of degree $n \geq 5$, satisfying $p(-i) = p(i)$, then there is at least one critical point of p in the union $D[-c; r] \cup D[c; r]$ of two closed disks with centres $-c, c$ and radius r , where

$$c = \cot(2\pi/n), \quad r = 1/\sin(2\pi/n).$$

If $n = 3$, then the closed disk $D[0; 1/\sqrt{3}]$ has this property; and if $n = 4$ then the union of the closed disks $D[-1/3; 2/3] \cup D[1/3; 2/3]$ has this property. In the last two cases, the domains are minimal, with respect to inclusion, having this property.

This theorem is stronger than any other known Rolle's Theorem for complex polynomials of any degree. A minimal Rolle's domain are found for polynomials of degree 3 and 4, answering Tschakaloff's question.

This is a joint work with Blagovest Sendov from the Bulgarian Academy of Sciences.